

# EXPLICIT SPECTRUM OF A CIRCULANT-TRIDIAGONAL MATRIX WITH APPLICATIONS

EMRAH KILIÇ AND AYNUR YALÇINER

ABSTRACT. We consider a circulant-tridiagonal matrix and compute its determinant by using generating function method. Then we explicitly determine its spectrum. Finally we present applications of our results for trigonometric factorizations of the generalized Lucas sequences.

## 1. INTRODUCTION

Tridiagonal matrices have been used in many different fields, especially in applicative fields such as numerical analysis (e.g., orthogonal polynomials), engineering, telecommunication system analysis, system identification, signal processing (e.g., speech decoding, deconvolution), special functions, partial differential equations and naturally linear algebra (see [1, 3, 4, 12, 16, 17]). Some authors consider a general tridiagonal matrix of finite order and then compute its LU factorizations, determinant and inverse (see [2, 5, 8, 13]).

A tridiagonal Toeplitz matrix of order  $n$  has the form:

$$A_n = \begin{bmatrix} a & b & & & 0 \\ c & a & b & & \\ & c & a & & \\ & & \ddots & \ddots & b \\ 0 & & & c & a \end{bmatrix},$$

where  $a, b$  and  $c$ 's are nonzero complex numbers.

A tridiagonal 2-Toeplitz matrix has the form:

$$T_n = \begin{bmatrix} a_1 & b_1 & 0 & 0 & 0 & \cdots \\ c_1 & a_2 & b_2 & 0 & 0 & \cdots \\ 0 & c_2 & a_1 & b_1 & 0 & \cdots \\ 0 & 0 & c_1 & a_2 & b_2 & \cdots \\ 0 & 0 & 0 & c_2 & a_1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix},$$

where  $a, b$  and  $c$ 's are nonzero complex numbers.

Let  $a_1, a_2, b_1$  and  $b_2$  be real numbers. The period two second order linear recurrence system is defined to be the sequence  $f_0 = 1$ ,  $f_1 = a_1$ , and

$$f_{2n} = a_2 f_{2n-1} + b_1 f_{2n-2} \quad \text{and} \quad f_{2n+1} = a_1 f_{2n} + b_2 f_{2n-1}$$

---

2000 *Mathematics Subject Classification.* 15A15, 15B36, 65Q05.

*Key words and phrases.* Circulant and tridiagonal matrices, spectrum, determinant, generating function.

for  $n \geq 1$ . Let  $D = a_1a_2 + b_1 + b_2$  and  $D^2 - 4b_1b_2 \neq 0$ .

Gover [6] and Marcellán and Petronilho [15] showed that the eigenvalues of matrix  $T_{2n+1}$  are  $a_1$  and

$$\frac{a_1 + a_2}{2} \pm \sqrt{\left(\frac{a_1 - a_2}{2}\right)^2 + b_1c_1 + b_2c_2 + 2\sqrt{b_1b_2c_1c_2} \cos \frac{k\pi}{n+1}}$$

for  $1 \leq k \leq n$ .

They also gave a closed equation for the eigenvalues of  $T_{2n}$ : They are the solutions of the following quadratic equations

$$(\lambda - a_1)(\lambda - a_2) - \left[b_1c_1 + b_2c_2 + \sqrt{b_1b_2c_1c_2}z_{nk}\right] = 0, \quad k = 1, 2, \dots, n,$$

where  $z_{nk}$ ,  $k = 1, 2, \dots, n$ , are the zeros of the polynomial  $R_n(z)$  defined by

$$R_{n+1}(x) = xR_n(x) - R_{n-1}(x), \quad n \geq 1$$

with initials  $R_0(x) = 1$ ,  $R_1(x) = x + \beta$  where  $\beta^2 = b_2c_2/b_1c_1$ .

Meanwhile, a matrix  $C_n$  is called a circulant matrix if it has the form

$$C_n = \begin{bmatrix} a_1 & a_2 & a_3 & \dots & a_{n-1} & a_n \\ a_n & a_1 & a_2 & \dots & a_{n-2} & a_{n-1} \\ a_{n-1} & a_n & a_1 & \dots & a_{n-3} & a_{n-2} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ a_3 & a_4 & a_5 & \dots & a_1 & a_2 \\ a_2 & a_3 & a_4 & \dots & a_n & a_1 \end{bmatrix}.$$

Circulant matrices are a special type of Toeplitz matrix and have many interesting properties. Circulant matrices have been used in many areas such as physics, differential equations and digital image processing. Also circulant and skew circulant matrices have become an important tool in networks engineering.

Define generalized Fibonacci and Lucas sequences by the recursion for  $n > 1$

$$U_n = PU_{n-1} - QU_{n-2} \quad \text{and} \quad V_n = PV_{n-1} - QV_{n-2},$$

with the initials  $U_0 = 0, U_1 = 1$ , and,  $V_0 = 2, V_1 = P$ , resp. When  $P = 1$  and  $Q = -1$ ,  $U_n = F_n$  ( $n$ th Fibonacci number) and  $V_n = L_n$  ( $n$ th Lucas number).

Recently some authors have studied various interesting combinatorial matrices defined by terms of certain sequences. We could refer to the works [10, 14, 11, 18] for details about combinatorial matrix examples: In [9], the authors consider skew circulant type matrices with any continuous Fibonacci numbers. Then they discuss the invertibility of the skew circulant type matrices and present explicit determinants and inverse matrices of them.

In this paper, we consider a circulant-tridiagonal matrix and then compute its determinant by using generating function method. Then we explicitly determine its all eigenvalues. We show that determinant of the matrix satisfies a period second order recurrence system. We give applications for trigonometric factorizations of the Lucas sequences.

## 2. THE MAIN RESULTS

We define a tridiagonal matrix  $H_n = [h_{ij}]$  of order  $n$  with  $h_{11} = a$ ,  $h_{i+1,i} = h_{i,i+1} = b$  for odd  $i$ ,  $h_{i,i+1} = h_{i+1,i} = a$  for even  $i$  and  $h_{nn} = a$  if  $n$  is even and  $h_{nn} = b$  if  $n$  is odd.

Then the matrix  $H_{2n}$  takes the form

$$H_{2n} = \begin{bmatrix} a & b & & & & \\ b & 0 & a & & & \\ & a & 0 & b & & \\ & & b & \ddots & \ddots & \\ & & & \ddots & 0 & b \\ & & & & b & a \end{bmatrix}.$$

We define a period two second order linear recurrence system  $\{h_n\}$  given by  $h_0 = 0$ ,  $h_1 = a + b$ , and the recursions

$$\begin{aligned} h_{2n} &= (a - b)h_{2n-1} - abh_{2n-2}, \\ h_{2n+1} &= (-a + b)h_{2n} - abh_{2n-1} \end{aligned}$$

for  $n \geq 1$ .

We give relationships between the period two second order linear recurrence system  $\{h_n\}$  and the determinant of  $H_n$ . Then we determine the eigenvalues of matrix  $H_n$ .

By expanding the determinant of  $H_n$  with respect to the first row, we have the following result without proof.

**Lemma 1.** For  $n > 1$ ,

$$\det H_{2n} = (-1)^{n+1} (a^{2n} - b^{2n}) \text{ and } \det H_{2n+1} = (-1)^n (a^{2n+1} + b^{2n+1}).$$

Let

$$H(x) = \sum_{n \geq 0} h_n x^n.$$

Also let

$$H_1(x) = \sum_{n \geq 0} h_{2n+1} x^{2n+1} \text{ and } H_2(x) = \sum_{n \geq 0} h_{2n} x^{2n}.$$

By these equations, we get the equation system

$$\begin{aligned} H_2(x) &= (a - b)xH_1(x) - abx^2H_2(x), \\ H_1(x) - (a + b)x &= (-a + b)xH_2(x) - abx^2H_1(x). \end{aligned}$$

By Cramer solution of the system, we obtain

$$H_1(x) = \frac{(a + b)abx^3 + (a + b)x}{1 + (a^2 + b^2)x^2 + a^2b^2x^4}$$

and

$$H_2(x) = \frac{(a^2 - b^2)x^2}{1 + (a^2 + b^2)x^2 + a^2b^2x^4}.$$

Thus we get

$$H(x) = H_1(x) + H_2(x) = \frac{(b + a)x + (a^2 - b^2)x^2 + (a^2b + ab^2)x^3}{1 + (a^2 + b^2)x^2 + a^2b^2x^4}.$$

Here note that

$$\begin{aligned} 1 + (a^2 + b^2)x^2 + a^2b^2x^4 &= (1 - \alpha^2x^2)(1 - \beta^2x^2) \\ &= (1 - \alpha x)(1 + \alpha x)(1 - \beta x)(1 + \beta x), \end{aligned}$$

where  $\alpha = \mathbf{i}a$ ,  $\beta = \mathbf{i}b$  and  $\mathbf{i} = \sqrt{-1}$ .

By partial fraction decomposition, we find  $A_1$ ,  $A_2$ ,  $B_1$  and  $B_2$  such that

$$H(x) = \sum_{n \geq 0} h_n x^n = \frac{A_1}{(1 - \alpha x)} + \frac{A_2}{(1 + \alpha x)} + \frac{B_1}{(1 - \beta x)} + \frac{B_2}{(1 + \beta x)}.$$

Solving the equation above, we get the coefficients have the forms:

$$A_1 = -\frac{(1+i)}{2}, \quad A_2 = -\frac{1-i}{2}, \quad B_1 = \frac{1-i}{2} \quad \text{and} \quad B_2 = \frac{1+i}{2}.$$

Therefore

$$h_n = \frac{1}{2} \left( -(1+i)\alpha^n - (1-i)(-\alpha)^n + (1-i)\beta^n + (1+i)(-\beta)^n \right).$$

Especially we have that

$$h_{2n+1} = (-1)^n (a^{2n+1} + b^{2n+1}) \quad \text{and} \quad h_{2n} = (-1)^{n+1} (a^{2n} - b^{2n}).$$

By the results above, we have the following result:

**Corollary 1.** For  $n > 1$ ,

$$\det H_n = h_n.$$

If we choose  $a$  and  $b$  as the roots of the characteristic equation of the general Lucas sequences,  $x^2 - Px + Q = 0$ , then we have

$$h_{2n+1} = (-1)^n V_{2n+1} \quad \text{and} \quad h_{2n} = (-1)^{n+1} U_{2n} \sqrt{\Delta},$$

where  $\Delta = P^2 - 4Q$ .

**Lemma 2.** For  $n > 1$ ,  $\det H_n = 0$  if and only if

$$\begin{cases} a = \pm b \text{ or } a^2 + b^2 = 2ab \cos \frac{2k\pi}{n} & \text{for } 1 \leq k \leq \frac{n-2}{2} & \text{if } n \text{ is even,} \\ a + b = 2\sqrt{ab} \cos \frac{(2k-1)\pi}{2n} & \text{for } 1 \leq k \leq n & \text{if } n \text{ is odd.} \end{cases}$$

*Proof.* Since  $h_n = \det H_n$  and by our result mentioned before,  $h_n = \det H_n = 0$  if and only if

$$\frac{1}{2} \left( -(1+i)\alpha^n - (1-i)(-\alpha)^n + (1-i)\beta^n + (1+i)(-\beta)^n \right) = 0.$$

Thus

$$(1+i)\alpha^n + (1-i)(-1)^n \alpha^n = (1-i)\beta^n + (1+i)(-1)^n \beta^n$$

or

$$\frac{\alpha^n}{\beta^n} = \frac{(1-i) + (1+i)(-1)^n}{(1+i) + (1-i)(-1)^n}.$$

For even  $n$  such that  $n = 2m$ , we find the all solution of the equation

$$\frac{\alpha^{2m}}{\beta^{2m}} = \frac{(1-i) + (1+i)}{(1+i) + (1-i)} = 1,$$

which, by  $\alpha = ia$  and  $\beta = ib$ , satisfies

$$a^{2m} = b^{2m}$$

or

$$a^{2m} - b^{2m} = 0.$$

From (pp. 34, formula 1.396.2, [7]), we recall the known result

$$\prod_{k=1}^{m-1} \left( x^2 + 1 - 2x \cos \frac{k\pi}{m} \right) = \frac{x^{2m} - 1}{x^2 - 1}.$$

By taking  $x = a/b$ , we get

$$(a^2 - b^2) \prod_{k=1}^{m-1} \left( a^2 + b^2 - 2ab \cos \frac{k\pi}{m} \right) = a^{2m} - b^{2m}.$$

Thus  $a^{2m} - b^{2m} = 0$  if and only if

$$a = \pm b \text{ or } a^2 + b^2 = 2ab \cos \frac{k\pi}{m}$$

for some  $1 \leq k \leq m-1$  where  $n = 2m$ .

For odd  $n$  such that  $n = 2m+1$ ,  $\det H_n = 0$  if and only if

$$\frac{\alpha^{2m+1}}{\beta^{2m+1}} = -1$$

or

$$\alpha^{2m+1} + \beta^{2m+1} = 0,$$

which, by  $\alpha = ia$  and  $\beta = ib$ , is equivalent to

$$a^{2m+1} + b^{2m+1} = 0.$$

By the product form of Chebyshev polynomials of the first kind, we have that

$$a^m + b^m = \prod_{k=1}^m \left( (a+b) - 2\sqrt{ab} \cos \frac{(2k-1)\pi}{2m} \right)$$

and so

$$a^{2m+1} + b^{2m+1} = \prod_{k=1}^{2m+1} \left( (a+b) - 2\sqrt{ab} \cos \frac{(2k-1)\pi}{2(2m+1)} \right).$$

Finally  $a^{2m+1} + b^{2m+1} = 0$  if and only if  $a+b = 2\sqrt{ab} \cos \frac{(2k-1)\pi}{2(2m+1)}$  for  $1 \leq k \leq 2m+1$ . Thus the claim is proven.  $\square$

Now we can determine eigenvalues of the matrix  $H_n$ . For this, we define a new period second order recurrence system  $\{g_n\}$  by the recursion

$$\begin{aligned} g_{2n} &= (a-b-x)g_{2n-1} - (ab)g_{2n-2}, \\ g_{2n+1} &= (-a+b-x)g_{2n} - (ab)g_{2n-1} \end{aligned}$$

with  $g_0 = 0$  and  $g_1 = a+b-x$  for  $n > 0$ .

On the other hand, by straightforward computations gives us the characteristic equation of matrix  $H_n$  as

$$\det(H_{2n} - xI_{2n}) = g_{2n}$$

and

$$\det(H_{2n+1} - xI_{2n+1}) = g_{2n+1},$$

where  $I_n$  is the identity matrix of order  $n$ .

Combining them, we determine the eigenvalues of matrix  $H_n$  :

**Theorem 1.** *The eigenvalues of  $H_{2n}$  are*

$$a \pm b \text{ and } \pm \sqrt{a^2 + b^2 - 2ab \cos \frac{k\pi}{n}}, \quad 1 \leq k \leq n-1,$$

and the eigenvalues of  $H_{2n+1}$  are

$$a + b \text{ and } \pm \sqrt{(a^2 + b^2) - 2ab \cos \frac{(2k-1)\pi}{2n+1}}, \quad 1 \leq k \leq n.$$

### 3. TWO APPLICATIONS

Now we will give two applications of our results on trigonometric factorizations of the second order recurrences  $\{U_n\}$  and  $\{V_n\}$ . If we choose the entries of the matrix  $H_n$  as  $a = (P + \sqrt{P^2 - 4Q})/2$  and  $b = (P - \sqrt{P^2 - 4Q})/2$ , then we will obtain the following results :

$$V_{2n+1} = V_1 \prod_{k=1}^n \left( V_2 - 2Q \cos \frac{2k-1}{2n+1} \pi \right)$$

and

$$U_{2n} = U_2 \prod_{k=1}^{n-1} \left( V_2 - 2Q \cos \frac{k\pi}{n} \right).$$

Especially when  $P = 1$  and  $Q = -1$ , then we get

$$L_{2n+1} = \prod_{k=1}^n \left( 3 + 2 \cos \frac{2k-1}{2n+1} \pi \right) \text{ and } F_{2n} = \prod_{k=1}^{n-1} \left( 3 + 2 \cos \frac{k\pi}{n} \right).$$

### REFERENCES

- [1] A. Bunse-Gerstner, R. Byers and V. Mehrmann, A chart of numerical methods for structured eigenvalue problems, *SIAM J. Matrix Anal. Appl.* 13(2) (1992), 419–453, doi: 10.1137/0613028.
- [2] R.L. Burden, J.D. Fairs and A.C. Reynolds, *Numerical Analysis*, 2nd Ed., Prindle, Weber & Schmidt, Boston, MA, 1981.
- [3] W. Chu, Fibonacci polynomials and Sylvester determinant of tridiagonal matrix. *Applied Math. Comput.* 216(3) (2010), 1018–1023, doi: 10.1016/j.amc.2010.01.089.
- [4] C.F. Fischer and R.A. Usmani, Properties of some tridiagonal matrices and their application to boundary value problems, *SIAM J. Numer. Anal.* 6(1) (1969), 127–142, doi: 10.1137/0706014.
- [5] C. M. Fonseca, On the eigenvalues of some tridiagonal matrices, *J. Comput. Appl. Math.* 200(1) (2007), 283–286, doi: 10.1016/j.cam.2005.08.047.
- [6] M.J.C. Gover, The eigenproblem of a tridiagonal 2-Toeplitz Matrix, *Linear Algebra Appl.* 197-198 (1994), 63–78, doi: 10.1016/0024-3795(94)90481-2.
- [7] I.S. Gradshteyn and I.M. Ryzhik, *Table of Integrals, Series, and Products*, Academic Press, San Diego, 1980.
- [8] W.W. Hager, *Applied Numerical Linear Algebra*, Prentice-Hall Inter. Editions, Englewood Cliffs, NJ, 1988.
- [9] Z. Jiang, Y. Jinjiang and F. Lu, On skew circulant type matrices involving any continuous Fibonacci numbers, *Abstract and Applied Analysis*, Article ID 483021, 10 pp., 2014, doi: 10.1155/2014/483021.
- [10] Z. Jiang and D. Li, The invertibility, explicit determinants, and inverses of circulant and left circulant and  $g$ -circulant matrices involving any continuous Fibonacci and Lucas numbers, *Abstr. Appl. Anal.* Article ID 931451, 14 pp., 2014, doi: 10.1155/2014/931451.
- [11] Z. Jiang, H. Xin and F. Lu, Gaussian Fibonacci circulant type matrices, *Abstr. Appl. Anal.* Article ID 592782, 10 pp., 2014, doi: 10.1155/2014/592782.
- [12] E. Kiliç and D. Tasci, Factorizations and representations of the backward second-order linear recurrences, *J. Comput. Appl. Math.* 201(1) (2007), 182–197, doi: 10.1016/j.cam.2006.02.012.
- [13] D.H. Lehmer, Fibonacci and related sequences in periodic tridiagonal matrices, *The Fibonacci Quarterly* 13(2) (1975), 150–158.

- [14] T. Mansour, A formula for the generating functions of powers of Horadam's sequence, Australasian Journal of Combinatorics 30 (2004), 207–212.
- [15] F. Marcellán and J. Petronilho, Eigenproblems for tridiagonal 2-Toeplitz matrices and quadratic polynomial mappings, Linear Algebra Appl. 260 (1997), 169–208, doi: 10.1016/S0024-3795(97)80009-2.
- [16] M. El-Mikkawy and E-D. Rahmo, A new recursive algorithm for inverting general tridiagonal and anti-tridiagonal matrices. Appl. Math. Comput. 204(1) (2008), 368–372, doi: 10.1016/j.amc.2008.06.053.
- [17] M. El-Mikkawy and F. Atlan, A new recursive algorithm for inverting general  $k$ -tridiagonal matrices. Appl. Math. Lett. 44 (2015), 34–39, doi: 10.1016/j.aml.2014.12.018.
- [18] J. Zhou and Z. Jiang, Spectral norms of circulant-type matrices with binomial coefficients and harmonic numbers, Int. J. Comput. Methods 11(5) (2014), 1-14, doi: 10.1142/S021987621350076X.

TOBB UNIVERSITY OF ECONOMICS AND TECHNOLOGY MATHEMATICS DEPARTMENT 06560 ANKARA  
TURKEY

*E-mail address:* `ekilic@etu.edu.tr`

SELÇUK UNIVERSITY SCIENCE FACULTY MATHEMATICS DEPARTMENT 42075 KONYA TURKEY

*E-mail address:* `ayalciner@selcuk.edu.tr`